Cytokine Test Requisition Form

Clear Form





See Page 2 for a list of Diagnostic Tests. Please mark ALL tests to be done. Patient Information, Referring Physician, and Ordering Laboratory are ALL REQUIRED for samples to be processed without delay.

PATIENT INFORMATION	REFERRING PHYSICIAN INFORMATION
Patient Name Surname: First Name: Sex Female Male Personal Health Number Birth (dd/mm/yy)	Physician Name (Surname, First name) Phone Number Fax Number Email address Comments
ORDERING LABORATORY INFORMATION	SAMPLE INFORMATION
Laboratory Name Address	Sample Type Serum Plasma CSF Time and Date Collected (dd/mm/yy) Ordering Lab Reference # Reason for Testing / Comments
Phone number Fax number Email address	Sample Attestation I attest that the sample was prepared in accordance with the sample collection procedure below

Sample Collection and Preparation Procedure: Collect blood in a plasma EDTA tube (Purple Vacutainer) or in a serum collection tube (Gold (SST) or Red Vacutainer) - please do not use heparinized tubes for sample collection. Within 30 minutes from collection, centrifuge at 1000 x g for 10 minutes at 4°C. Immediately transfer/aliquot 0.3 - 1 mL cell-free plasma or serum to a small tube (~3mL or smaller tube such as a false-bottom tube). Freeze the sample (≤ -20°C) and ship on dry ice (consider including a temperature monitoring device in the shipment).

SHIPPING INFORMATION

Please ship on Monday or Tuesday of any desired workweek to ensure your samples are scheduled to arrive before 3pm on Friday. Please do not send samples on weekends or holidays as Eve Diagnostics will be closed and samples will not be received until the following workweek. Shipments from any shipping company are accepted (including Purolator and FedEx). Tracking numbers should be retained for delivery confirmation. If problems arise or your package is not delivered promptly please email us at contact@evetechnologies.com.

Shipping -- See Sample Shipping Guide. Please send properly labeled and packaged samples with this requisition to: Eve Technologies Corporation, 3415 A 3rd Avenue NW, Calgary, AB, Canada, T2N 0M4

RECEIVING RESULTS

Results typically follow within 5 business days after receipt of sample and payment.

Results reports will be attached to your online order. If requested, reports can also be faxed to referring physicians/laboratories - please provide a valid fax number to receive faxed results. If you have not received your results, please contact the laboratory that sent your sample. If you have not received your report, contact the laboratory that processed your sample. For missing or delayed reports, please reach out to us.

CONTACT INFO FOR LABS

Eve Diagnostics

3415A 3rd Ave NW Calgary, AB, T2N 0M4 Phone: 587-975-8850 Fax: 4587-975-8899 Email: contact@evetechnologies.com

Visit our website: www.evetechnologies.com/cytokine-array-patient-diagnostics/

Cytokine Test Requisition Form





Eve Technologies is certified by Centers for Medicare & Medicaid Services (CMS) as a High Complexity International Laboratory under the Clinical Laboratory Improvement Amendments (CLIA); speciality Diagnostic Immunology, subspeciality General Immunology. These are Laboratory Developed Tests (LDT) and do not appear on the lists of tests in the Federal Register and have not been reviewed by the U.S. Food and Drug Administration.

	IMMUNE BIOMARKER TEST REQUISITION
	Medical Personnel: Please mark <u>ALL</u> tests to be done
	Cytokine, Chemokine, Growth Factor Panel (71-Plex): 6Ckine, BCA-1, CTACK, EGF, ENA-78, Eotaxin, Eotaxin-2, Eotaxin-3, FGF-2, Flt-3 Ligand, Fractalkine, G-CSF, GM-CSF, GROα, I-309, IFNα2, IFNγ, IL-1α, IL-1β, IL-1β, IL-1β, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12(p40), IL-12(p70), IL-13, IL-16, IL-17A, IL-17E/IL-25, IL-17F, IL-18, IL-20, IL-21, IL-22, IL-23, IL-27, IL-29A, IL-33, IP-10, LIF, MCP-1, MCP-2, MCP-3, MCP-4, M-CSF, MDC, MIG, MIP-1α, MIP-1β, MIP-1δ, PDGF-AA, PDGF-AB/BB, RANTES, sCD40L, SCF, SDF-1α+β, TARC, TGFα, TNFα, TNFβ, TPO, TRAIL, TSLP, VEGF-A Indications: Use this test when identifying relevant therapy targets, mechanism of disease is unknown, arthritis, macrophage activation syndrome (MAS), Hemophagocytic lymphohisticcytosis (HLH), sepsis / toxic shock, necrotizing fasciitis, admitted meningitis, admitted pneumonia, Kawasaki syndrome, prolonged or periodic fever, severe or chronic inflammation, organ dysfunction. Also useful in investigation of 'cytokine storm' seen in HLH and related conditions. Reference Intervals available for PLASMA-EDTA samples, SERUM, and CSF samples Results presented with cytokine groupings (immune signatures)
	Focused Cytokine, Chemokine, Growth Factor Panel (15-Plex): GM-CSF, IFNy, IL-1β, IL-1RA, IL-2, IL-4, IL-5, IL-6, IL-8, IL-10, IL-12(p40),
L	IL-12(p70), IL-13, MCP-1, TNF-α
	Indications: Use this test when identifying relevant therapy targets in severe or chronic inflammation, and 'cytokine storm' seen in COVID-19. Reference Intervals available for PLASMA-EDTA samples, SERUM, and CSF samples
ſ	Soluble Cytokine Receptor Panel (14-Plex): sCD30, sEGFR, sgp130, slL-1Rl, slL-1Rll, slL-2Ra, slL-4R, slL-6R, sRAGE, sTNFRl, sTNFRll,
L	sVEGFR1, sVEGFR2, sVEGFR3
	Indications: Use this test when identifying relevant therapy targets, mechanism of disease is unknown, arthritis, macrophage activation syndrome (MAS), Hemophagocytic lymphohisticcytosis (HLH), sepsis / toxic shock, necrotizing fasciitis, admitted meningitis, admitted pneumonia, Kawasaki syndrome, prolonged or periodic fever, severe or chronic inflammation, organ dysfunction. Reference Intervals available for PLASMA-EDTA samples and CSF samples
ſ	Serum Amyloid A (SAA) / ADAMTS13 Panel (2 Plex)
	Indications: Use this test for acute phase inflammatory detection, chronic inflammatory diseases. Reference Intervals available for PLASMA-EDTA samples and CSF (SAA only) samples
	Complement Profile Panel (13-Plex): Adipsin (Factor D), C1q, C2, C3, C3b/iC3b, C4, C4b, C5, C5a, Factor B, Factor H, Factor I, Mannose-Binding Lectin Indications: Use this test for acute phase inflammatory detection. Reference Intervals available for PLASMA-EDTA samples.
Ī	Interferon Panel (9-Plex): IFNα2, IFNβ, IFNγ, IFNγR1, IFNε, IFNλ1, IFNλ2, IFNλ3, IFNω
	Indications: Use this test to assess the interferon response in viral infections, autoinflammatory interferopathies, and autoimmune diseases. Reference Intervals available for PLASMA-EDTA samples and SERUM samples.
Ī	Angiogenesis Panel (12-Plex): Angiopoietin-2, BMP-9, Endoglin, Endothelin-1, FGF-1, Follistatin, HB-EGF, HGF, Leptin, PLGF, VEGF-C, VEGF-D
L	Indications: Use this test to assess abnormalities of angiogenesis in cancer, autoimmune diseases, cardiovascular disease, or other chronic
	inflammatory conditions. Reference Intervals available for PLASMA-EDTA samples.
	SAA / Cardiovascular Disease Panel (6-Plex): ADAMTS13, D-Dimer, SAA, slCAM-1, sP-Selectin, sVCAM-1
	Indications: Use this test to investigate the underlying molecular basis of suspected coagulopathies, cardiovascular disease, and acute or chronic inflammation. Reference Intervals available for PLASMA-EDTA samples.
Ī	Long COVID Vascular Health Panel (18-Plex): ADAMTS13, Angiopoietin-2, BMP-9, D-Dimer, Endoglin, Endothelin-1, FGF-1, Follistatin,
	HB-EGF, HGF, Leptin, PLGF, SAA, slCAM-1, sP-Selectin, sVCAM-1, VEGF-C, VEGF-D Indications: Use this panel to investigate dysfunctions in vascular and endothelial integrity, coagulation, and inflammation. The panel consists of circulating factors that may suggest abnormalities in angiogenesis (a response to persistent vascular injury), coagulation, vascular function and inflammatory responses, which are thought to be key features of long COVID pathogenesis. Reference Intervals available for PLASMA-EDTA samples.
	COMMENTS



PHN